




Ivermectin BP 60mg
Praziquantel BP 375mg

DESCRIPTION

For the control of infestations due to adult and immature roundworms, lungworms and bots, with the addition of praziquantel to control tapeworm infestations:

- Ascarids (adults and 3rd and 4th stage larvae)
- Bots (oral and gastric larval stages)
- Hairworms (adults)
- Intestinal threadworms (adults)
- Large strongyles (adults)
- Strongylus vulgaris
- Large-mouth stomach worms (adults)
- Lungworms (adults and 4th stage larvae)
- Pinworms (adults and 4th stage larvae)
- Small strongyles (adults, and some resistant to class of benzimidazoles)
- Small strongyles (4th stage larvae)
- Summer sores and dermatitis caused by neck threadworm
- Tapeworm (anoplocephala perfoliate)



COMPOSITION

Each 9g sachet of film-coated granules contains:

- 60mg Ivermectin BP
- 375mg Praziquantel BP

- Ivermectin treatment dosage of 0.2mg/kg (0.09mg/lb) of horse's bodyweight
- Praziquantel treatment dosage of 1.5mg/kg (0.68mg/lb) of horse's bodyweight

HOW MANY DO I NEED?

Recommended Control Program

- ONE SACHET per 300kg (660lb)

IMPORTANT:

- Consult your veterinarian for a control program to meet your specific needs.
- Regular treatment will reduce the chances of colic caused by anoplocephala perfoliata and verminous arteritis caused by strongylus vulgaris.

OVERDOSAGE

Should a massive overdose occur, treat symptomatically and supportively

SIDE EFFECTS

Horses carrying heavy infections of onchocera microfilariae may have reactions such as swelling and itching after treatment. It is assumed that these reactions are the result of destruction of large numbers of microflora; consult your veterinarian if these signs persist.

WARNINGS

- Ablver Plus has been formulated specifically for use in horses. Ensure other animals do not have access to ingest the treatment, as they may be adversely affected by the concentration of ivermectin.
- Pregnant mares should be treated as normal during their pregnancy taking care not to stress them during treatment. However, in general, anthelmintics should not be administered to mares during the first 60 days of gestation. Pregnant mares should be administered ivermectin or praziquantel 1–3 days before foaling to prevent lactogenic transmission of strongyloides westeri, which can cause diarrhea in young foals.
- Do not treat horses intended for human consumption.
- Keep out of reach of children.
- Veterinary use only.

EXAMPLE DOSAGE CALCULATOR

Treatment Regime

Horse's Weight	Dosage Calculation
550kg (1210lbs)	1.66 sachets (1 measure for every 300kg/660lbs of bodyweight)

FEEDING TIPS

- Sprinkle directly onto feed.
- Combine with molasses or corn syrup and add to feed or treat (ie. apple).

INTERACTIONS / CONTRAINDICATIONS

Do not use in foals under 4 weeks of age.

TOXICITY

Reports indicate the combination of ivermectin and the silver nightshade plant results in a severe toxic reaction in horses.

DISPOSAL

- Dispose of unused product and containers in accordance with guidance from your local waste regulation authority.
- Do not contaminate water ways with the product or used containers.

STORAGE

- Store in a cool dry atmosphere away from extreme heat.
- Store/Transport at and 20° to 30°C (68° to 86°F)